CLAIMS

- 1. A process for producing 1-hexene, which comprises trimerizing ethylene in the presence of a catalyst comprising a tantalum compound and an alkylating agent containing a metal under a condition wherein the water content in the reaction system is 4 moles or less per mole of the tantalum atom, and/or under a condition wherein the amount of molecular oxygen in the reaction system is 2 moles or less per mole of the tantalum atom.
- The process for producing 1-hexene according to claim
 wherein the water content in the reaction system is 3 moles
 or less per mole of the tantalum atom.
 - 3. The process for producing 1-hexene according to claim 1, wherein the water content in the reaction system is 2 moles or less per mole of the tantalum atom.

15

25

- The process for producing 1-hexene according to claim
 wherein the water content in the reaction system is
 substantially anhydrous.
- 5. The process for producing 1-hexene according to any one of claims 1 to 4, wherein the amount of the molecular oxygen in the reaction system is 1.5 moles or less per mol of the tantalum atom.
 - 6. The process for producing 1-hexene according to any one of claims 1 to 4, wherein the amount of the molecular oxygen in the reaction system is 1 mol or less per mole of the tantalum atom.
 - 7. The process for producing 1-hexene according to any one of claims 1 to 4, wherein the amount of the molecular oxygen in the reaction system is substantially oxygen-free.

- 8. The process for producing 1-hexene according to any one of claims 1 to 7, wherein the tantalum compound is a tantalum halide.
- 9. The process for producing 1-hexene according to any one of claims 1 to 7, wherein the tantalum compound is tantalum pentachloride or tantalum pentabromide.

5

- 10. The process for producing 1-hexene according to any one of claims 1 to 9, wherein the alkylating agent is a hydrocarbyl metal, a hydrocarbyl metal halide, or an alkylaluminoxane.
- 11. The process for producing 1-hexene according to any one of claims 1 to 9, wherein the alkylating agent is tetramethyltin, tetraethyltin, dimethylzinc, methyllithium, trimethylaluminum, n-butyllithium, allyltriphenyltin, triethylaluminum, dimethylaluminum chloride, tetraphenyltin, methylaluminoxane, or methylmagnesium bromide.